




children at risk



How Air Pollution from Power Plants Threatens the Health of America's Children



MIKE SEAMANS

Prepared by



77 Summer Street
Boston, MA 02110

April 2002





Credits

Written by:

L. Bruce Hill, Ph.D. and Martha Keating,
Clean Air Task Force

Foreword by:

George Thurston, Sc.D.,
New York University School of Medicine

Design by:

Jill Bock Design

Printed by:

Spectrum Printing & Graphics

Acknowledgements

Children at Risk was prepared by the Clean Air Task Force for Clear the Air. Conrad Schneider, Clean Air Task Force, and Angela Ledford and Jamie Linski, Clear the Air, and Karen Hopfl-Harris, Physicians for Social Responsibility provided editorial comments. Accompanying fact sheets were developed by David Schoengold, MSB Energy Associates. Deborah Shprentz, Atmospherix, provided assistance with reference compilation.

This report was made possible with funding from The Pew Charitable Trusts. The opinions expressed in this report are those of the authors and do not necessarily reflect the views of The Pew Charitable Trusts.

Copies of the report and State Fact Sheets are available at:

<http://www.cleartheair.org>

<http://clnatf.org>



Foreword | April 2002

By George D. Thurston, Sc.D.
New York University School of Medicine

Millions of children in America today are exposed to unhealthy air at home, at school, or at their playground. Scores of new studies each year demonstrate that children are more susceptible to air pollution than adults. Studies indicate that exposure to air pollutants such as particulate matter, sulfate, sulfur dioxide gas, and ozone can result in reduced lung function, asthma attacks, increased visits to the doctors office and emergency rooms, hospitalizations and may, very tragically, also lead to increased risk of infant death.

Several factors may increase the risk of all children to air pollution relative to adults. One of the greatest causes is the higher activity level of children. Pound for pound, children breathe more air for their size than adults do. Children spend more time playing outdoors, which increases their exposure to outdoor air pollution. The lung's defense systems in children are still developing, and are thus unable to defend against the effects of pollutants as effectively as adult lungs. Children also suffer a higher prevalence of asthma than adults, and asthma makes kids far more susceptible to impacts of air pollution. Finally, a higher percentage of children than adults live in poverty, meaning that their access to health care is more limited, and recent studies indicate that air pollution affects those living in poverty more than those with means.

Health researchers have long known that air pollution reduces the lung function of children and causes asthma attacks, based on research conducted at schools and summer camps over the past few decades. Moreover, asthma has been on the rise in the U.S., having nearly doubled in the past two decades. Why is this? Is it, in part, due to some form of air pollution? We don't yet know. One California study suggests that kids who play sports year-round in polluted areas have more newly diagnosed cases of asthma. Another indicates that people who grow up in high ozone areas have a higher prevalence of asthma.

Children at Risk highlights recent research and describes links between pollutants associated with power plants and children's health. Studies across the world have linked particulate matter exposures to infant deaths. Moreover there is a suggested link between air pollution and adverse birth outcomes, such as slowed development and low birth weight in fetuses, coupled with higher premature births. Newborns also face setbacks from power plant pollutants and possible stunted lung development. All of these adverse outcomes put America's children at risk for health problems later in life.

Aging power plants are the chief sources of many of the pollutants that affect children in the U.S. For example, two thirds of the sulfur dioxide gas emitted in the U.S. comes from power plants. Sulfur dioxide, itself a potential health risk near smokestacks, converts into harmful sulfate particulate matter and sulfuric acid downwind of the plant.

Global warming, driven by our dependence on fossil fuels to generate electricity, presents different risks to children. In a recent health effects analysis¹, my co-investigators and I found that substantial public health gains will result in the nations that mitigate carbon dioxide emissions by switching from carbon intensive energy sources to cleaner technologies due to the associated reductions in particulate matter and ozone smog. The primary beneficiaries of these policies will be children.

In summary, numerous risk analyses have linked power plants to pollutants that can harm children. Considering these potential health risks, Congress should take action now to provide relief to our children by closing the Clean Air Act loophole that still allows hundreds of power plants to avoid modern pollution standards some 30 years after the Act was made law and by requiring steep cuts in mercury and carbon dioxide emissions.



George D. Thurston, Sc.D.
New York University, April 2002

Executive Summary

Whether at home, school, or play, children are exposed to emissions from power plants. This report reviews important recent advances in our understanding of the link between air pollution and children's health. A number of harmful pollutants are emitted by power plants. Thus, cleaning up power plants will have a great impact on the quality of children's health in America. This report can serve as an educational tool and an aide to healthcare providers, politicians and citizens who want to take action to protect children's health using the best science available. Cleaner air means healthier kids today and healthier adults tomorrow.

As this report shows, children are the most susceptible members of our society to the detrimental effects posed by air pollutants emitted from power plants. While countless studies have looked at the effects of various air pollutants on human health, few have focused specifically on children. However, children represent our future and to protect children is to protect all people of all ages. Children remain particularly susceptible to pollution because their defense mechanisms have not yet fully developed, increasing their susceptibility to the harmful effects of pollution. Children also breathe more rapidly and have more lung surface area for their body size compared to adults, which means they take in more air per minute and inhale more air for their size. In fact, pound-for-pound, children breathe 50 percent more air than do adults, and as a result, our children inhale a greater percentage of pollution. Children also spend more time outdoors thus increasing their exposure to outdoor air pollutants. Because exercise increases the penetration of pollutants into the lungs, our children's outdoor activities make adverse health effects more likely. This is of particular concern because tens of thousands of schools are located near the most polluting, outdated power plants.

Power plant emissions and their byproducts form particulate matter, ozone smog and air toxics. These pollutants are associated with respiratory hospitalizations, lost school days due to asthma attacks, low birth weight, stunted lung growth and tragically, even infant death. Air pollution is a pervasive problem across America for urban, suburban, and rural communities. It is an inescapable fact that air pollution is everywhere – indoors and out – and kids breathe and absorb more of it than adults do.

Our children are at risk from power plant pollution:

- Over 25 million children in the U. S. live in counties that violate national air quality standards for the common pollutants ozone, particulate matter and sulfur dioxide;
- Cases of asthma have rapidly increased, more than doubling in the past two decades. Six percent of U.S. children have asthma;
- Thirty-five million of our children live within 30 miles of a power plant—a distance within which local communities may reasonably be affected by a power plant's smoke plume; an estimated 2 million of these children are asthmatic and are particularly susceptible to these pollutants;
- 72,000 of our schools are within 30 miles of a power plant;
- Average health risks to children due to exposure to power plant combustion wastes could be up to 10,000 times higher than EPA's allowable risk levels for cancer and other illnesses.

Power plants are a major source of the most common pollutants in the air that harm children. Power plants emit 67 percent of the sulfur dioxide (SO₂), 23 percent of the nitrogen oxides (NO_x), 33 percent of the mercury, and 38 percent of the carbon dioxide from energy related sources. In much of the U.S., especially in the East, Midwest and South, sulfates make up the bulk of so-called fine particulate matter. Power plants are responsible for about half of the fine particulate matter in many parts of the U.S. Numerous epidemiological studies have suggested that sulfate particles are among those most strongly associated with health impacts and premature mortality in adults.

Coal-fired power plants are also the largest U.S. source of air toxics. Based on an analysis of 1998 Toxics Release Inventory data, power plants ranked 5th in releases of developmental and neurological toxins with a total of 78 million pounds released to the air and surface waters.

Key findings of recent studies include:

For all children:

- As fine particulate levels rise, emergency room visits by asthmatic children also increase, even when fine particulate levels are below EPA's air quality standard;
- Exposure to particulate matter can slow lung function growth in children;
- Children living in high ozone communities and who played sports year-round were three times more likely to develop asthma compared to children who did not play sports. This is some of the first evidence suggesting smog can cause asthma;
- Methylmercury can have adverse effects on the developing and adult cardiovascular systems, blood pressure regulation and heart-rate variability;
- Global warming could lead to more frequent and severe air pollution problems, the spread of infectious and communicable diseases, and increasingly extreme weather events such as heat waves that could disproportionately affect children.

For unborn children (prenatal):

- A new California study suggests prenatal ozone exposures may cause heart defects;
- Research in a coal dominated region of the Czech Republic indicates that stunted development in unborn children may be a result of exposure to very high levels particulate matter;
- Researchers in China have found that high concentrations of particulate matter may affect developing babies;
- Methylmercury interferes with the development and function of the central nervous system. Prenatal exposure from maternal consumption of mercury-contaminated fish can result in problems later in childhood such as learning disabilities, attention deficits, loss of

IQ points or other disorders depending on the severity of exposure. Ten percent of women of child-bearing age are estimated to carry a body burden of mercury contamination above EPA's safe level.

For newborns:

- U.S. researchers in a study of 86 cities found that infants who lived in a highly polluted city during their first two months of life had a mortality rate ten percent higher than infants living in the city with the cleanest air;
- A preliminary study projects that eleven percent of the infant mortality in the United States is attributable to particulate matter even at low to moderate levels;
- Exposure to ozone may permanently affect lung structure of children; monkeys exposed to ozone developed little more than half of the normal number of branches of their lungs compared to monkeys exposed only to clean air;
- A recent study suggests that asthmatic children that were born pre-term and/or with low birth weights, are at greater risk from ozone exposures;
- The ten percent of women above EPA's safe level of mercury translates nationally into 6 million women of childbearing age with elevated levels of mercury from eating contaminated fish, and approximately 390,000 newborns at risk of neurological effects from being exposed in utero to elevated levels of mercury.

Association between Air Pollution and Lung Growth in Southern California Children

W. JAMES GAUDERMAN, ROB McCONNELL, FRANK GILLILAND, STEPHANIE LONDON, EDWARD AVOL, HITA VORA, KIROS BERHANE, EDWARD B. RAPPAPORT, FRED LURN, HELENE G. MARGOLIS, and JOHN PETERS
Department of Preventive Medicine, University of Southern California School of Medicine, Los Angeles; Air Resources Board, State of California, Sacramento, California; and National Institute of Environmental Health Sciences, Research Triangle Park, North Carolina

Fetal Growth and Maternal Exposure to Particulate Matter

Jan Dejmek,¹ Sherry G. Selevan,² Ivan Benes,³ Ivo Solansky,¹ and Rana...



American Journal of Epidemiology
Copyright © 2002 by the Johns Hopkins Bloomberg School of Public Health
All rights reserved.

Ambient Air Pollution and Risk of Birth Defects in Southern California

Beate Ritz,^{1,2} Fei Yu,³ Scott Fruin,^{4,5} Guadalupe Chapa,⁴ Gary M. Shaw,⁶ and John A. Hertz...

Association between Air Pollution and Low Birth Weight based Study

Xiaobin Wang,¹ Hui Ding,² Louise Ryan,³ and Xiping Xu^{4,5}
¹Department of Pediatrics, Boston University School of Medicine and Boston Children's Hospital, Boston, MA; ²Department of Biostatistics, Harvard School of Public Health, Boston, MA; ³Department of Environmental Health, Harvard School of Public Health, Boston, MA; ⁴Department of Environmental Health, Harvard School of Public Health, Boston, MA; ⁵Department of Environmental Health, Harvard School of Public Health, Boston, MA

Air Pollution and Exacerbation of Asthma in African American Children in Los Angeles

Bart Ostro,¹ Michael Lipscomb,^{1,2} Hazel Braxton-Owens,³ and Mary Whit...

The Effect of Ozone on Inner-City Children with Asthma: Identification of Susceptible Subgroups

KATHLEEN M. MORTIMER, IRA B. TAGER, DOUGLAS W. DOCKERY,
Department of Epidemiology and Environmental Health, Harvard School of Public Health, Boston, MA; Department of Epidemiology, University of California, Berkeley, California; Epidemiology and Biostatistics, University of California, Berkeley, California; Environmental Health Effects Research Laboratory, U.S. Environmental Protection Agency, Chapel Hill, NC; and ⁴Western Reserve University, Rainbow Babies and Children's Hospital, Cleveland, OH

Lung Function Growth and Ambient Ozone: A Three-Year Population Study in School Children

THOMAS FRISCHER, MICHAEL STUDNICKA, CHRISTIAN GARTNER, ERICH TAUBER, FRITZ ANDREAS VEITER, JOHN SPENGLER, JOACHIM KÜHR, and RADVAN URBANEK

Asthma in exercising children exposed to ozone: a population-based study

Rob McConnell, Kiros Berhane, Frank Gilliland, Stephanie J London, Talat Islam, W James Gauderman, C. M. Frisvold, John M. Peters, and John A. Hertz

Decline of Ambient Air Pollution and Respiratory Symptoms in Children

JOACHIM HEINRICH, BERND HOELSCHER, and Pulmologic Centre, First Internal Department, Vienna, Austria; Harvard School of Public Health, Boston, MA; and ³University Children's Hospital of Freiburg, Freiburg, Germany

Recommendations

While the benefits of reducing power plant pollution have been estimated for adults and are many times the cost of emissions controls, little work has been done to quantify the benefits for children. But, certainly the benefits will be great. Quality of life can be improved. Premature death can be avoided. The cost of health care can be decreased.

Comprehensively reducing pollution from coal-fired power plants will address each of the threats from air pollution that children face. Power plants must be required to comply with modern emission control standards. In addition, the nation's power fleet should be held to nationwide caps on all four of the key types of power plant pollution including nitrogen oxides, sulfur dioxide, mercury and other air toxics, and carbon dioxide. Reducing power plant emissions of nitrogen oxides and sulfur dioxide by at least 75 percent beyond current legal requirements will dramatically reduce fine particulate matter pollution so that children can breathe more easily.

The threat of power plant air pollution to children can only be meaningfully reduced when the Clean Air Act's 30-year loophole that allows old and dirty power plants to escape modern standards is finally closed. Once this is accomplished, U.S. energy policy will better account for

public health and the environmental costs associated with electricity production. This will propel us toward a more sustainable energy future that relies increasingly on cleaner sources of energy including renewable energy resources and conservation.

Strategies that will reduce carbon dioxide pollution from power plants will not only curb emissions of a greenhouse gas that causes climate change, but will provide the added benefits of reducing exposure to air pollutants, decreasing the risk of the spread of infectious diseases, and reducing temperature-related stress on children.

Our children's health and quality of life are harmed by air pollution today. The specter of global warming hangs over their future. We can leave our children a legacy of cleaner air and an improved environment by making wise choices today. Let's make comprehensive power plant clean up our gift to them.



Power Plant Emissions

A Threat to America's Children

Whether at home, school, or play, children are exposed to pollution in our air, land, and water from power plants. As this report shows, children are the most susceptible members of our community to the detrimental effects posed by these emissions. It is an inescapable fact that air pollution is everywhere – indoors and out – and kids breathe and absorb more of it than adults do. In addition to breathing harmful emissions from power plants, children are also exposed to contaminants in water and soil from disposal of coal combustion wastes.²

Air pollutants released by power plants are pervasive and harmful. They include particulate matter, ozone, sulfur dioxide gas, sulfate particulate matter, nitrogen oxides, mercury and a host of other air toxics. Collectively, these pollutants are associated with asthma attacks, respiratory disease, heart disease, and have been shown to retard cognitive development and stunt lung growth.^{3,4,5,6} A lifetime of exposure to many toxic air pollutants has been associated with cancer in adults. Exposure causes respiratory hospitalizations, lost school days due to asthma attacks, and is associated with low birth weight (birth weight is an important predictor of survival and illnesses in infants).⁷ In some cases, power plant pollutants may even cause death.^{8,9}

What can be done about the problem? Certainly technology is not a barrier to making progress in reducing the risks to our children. If today's modern smokestack emissions controls were used across the board, emissions of sulfur and nitrogen oxides (that form particulate matter and ozone smog) from old power plants would decrease by 90 percent or more.¹⁰ Combined use of sulfur dioxide and nitrogen oxides removal technologies can achieve substantial mercury reductions too. Committing to a more balanced energy policy will also help. An energy system that relies on conservation, clean renewables, and cleaner burning fossil technologies will reduce the amount of pollutants emitted and can reduce our dependence on carbon-intensive electric generating technologies which will help to stem the problem of global warming.

Currently, proposed state and federal legislation would significantly reduce the impact of power plant emissions on our children by cutting emissions nationally by 75 percent or more beyond the 1990 Clean Air Act Amendments and for the first time set emission standards for mercury and carbon dioxide. In addition to this legislation, it is also crucial that existing laws such as New Source Review, the National Ambient Air Quality

Standards and rules governing haze and hazardous air pollutants be maintained and fully enforced to prevent backsliding, as well as to ensure progress in children's respiratory health.

This report describes the risks to children posed by air pollution associated with coal-fired electricity generation. It summarizes the most recent research on how particulate matter, ozone, sulfur dioxide and hazardous air pollutants such as mercury affect children. Supplementing this report are state-by-state fact sheets with locations of power plants and statistics on the children at risk who live near them. These fact sheets can be found at – <http://www.cleartheair.org> or <http://clnatf.org>.



Children Face the Highest Risks

Air pollution is a pervasive problem across America for urban, suburban, and rural communities. Tens of thousands of schools are located near outdated, grandfathered power plants. Children are far more susceptible than adults to the adverse health effects of air pollution for a variety of reasons.^{11,12} First, children are more active than adults and therefore breathe more rapidly. Second, compared to adults, children (including teenagers) also have more lung surface area compared to their body weight that means that they inhale more air for their size. In fact, they breathe 50 percent more air pound-for-pound than adults do. As a result, our kids inhale a greater percentage of pollution for their size. Children also spend more time outdoors where air pollution may be higher. What's worse is that this combination of exercise and higher pollution levels outdoors increases the penetration of pollutants into the lungs, making children more likely to suffer adverse health effects. Moreover, children's defense mechanisms have not yet developed fully, which also increases their susceptibility to the harmful effects of pollution.

Children are Growing Up Breathing Dirty Air

National air quality standards were designed by Congress to protect all Americans from certain forms of common air pollution with "an adequate margin of safety", but our children are far from safe. Over 25 million

children in the U. S. live in counties that violate national air quality standards for the common pollutants ozone, particulate matter and sulfur dioxide.¹³ Nearly two million of the children in areas violating air standards are asthmatic and are particularly susceptible to these pollutants.



Many of America's children also live close to and go to school near power plants. Thirty-five million of our children live within 30 miles of a power plant – a distance within which local communities may be directly affected by a power plant's smoke plume.¹⁵ An estimated 2 million of these children are asthmatic. What's worse, 72,000 of our schools are within 30 miles of a power plant. (See state statistics, Appendix A and ranked exposure profiles, Appendix B.) Also, as described below, children living near power plants may also inhale various other air toxics emitted from the smokestack or may be exposed to pollutants in power plant combustion wastes released into ground water.

Power Plants are the Largest Sources of Air Pollution in the U.S.

Power plants are a major source of the most common pollutants in the air that harm children. In 1998, power plants emitted 67 percent of the sulfur dioxide (Figure 1) 25 percent of the nitrogen oxides, 34 percent of the mercury and 38 percent of the CO₂ in the United States.^{16,17,18} Moreover, after spewing from smokestacks right in our communities, these pollutants combine in

the atmosphere forming "secondary pollutants".¹⁹ Secondary pollutants, particularly ozone and sulfate, are some of the most harmful and widespread. For example, nitrogen oxides form acidic nitrate particulate matter, nitric acid droplets and ozone smog. Sulfur dioxide emissions from power plants form sulfate particulate matter and sulfuric acid. Sulfates are the dominant contributor to fine particulate matter in many areas of the U.S., especially in the eastern half.²⁰ Burning coal also releases highly toxic mercury and other toxic air emissions. Coal-fired power plants are the largest U.S. source of air toxics.²¹ Smokestacks directly emit toxic metals and gases directly into the atmosphere such as mercury, arsenic, chromium, beryllium and acid gases such as hydrochloric acid. Stack tests at the nation's coal-fired power plants have detected sixty-seven different air toxics emitted from the smokestacks.

Figure 1 – Sources of Sulfur Dioxide in the U.S., 1999.¹⁶ Electric utilities comprise two thirds of all sulfur emissions.



How Air Pollution Harms Children

The Children's Health Impacts Pyramid¹⁴

The health impacts of air pollution on children are illustrated in a conceptual pyramid diagram. The base of the pyramid represents the proportion of the population at risk from air pollution. Children represented along the base experience some symptoms beginning with subtle effects that may either go untreated or need no treatment. Of these children, however, some will have small changes in lung function, increased symptoms and medication use (higher up the pyramid). With more severe exposures, some children will see physicians and be prescribed medications or be admitted to the hospital. For the children represented by the top of the pyramid, in the worst instances, permanent changes may occur in the developing lungs of children and in some cases pollution can result in death.

Particulate Matter Dangerous for Children to Breathe

Particulate matter is, perhaps, the most pervasive and harmful pollutant from power plants plaguing America's children (Figure 2). During the hot, hazy days of summer, it is the particulate matter haze that you see hanging in the air not just humidity. While power plants directly emit some particulate matter as soot, the sulfur dioxide gas from power plants is a major source of particulate matter as it becomes transformed into tiny acidic sulfate particles in the atmosphere. These tiny particles are the most harmful and therefore of greatest concern. Fine particulate matter is of concern because it penetrates into our indoor living spaces thereby increasing our exposure. Fine particulate matter, known as PM_{2.5}, less than 2.5 microns in diameter or 1/100th the width of a human hair is deposited deep in the lung where it can affect both the respiratory and cardiovascular systems. Power plants release more tons of particulate matter-forming sulfur dioxide than any other pollution source. This means that power plants are responsible for about half of the fine particulate matter in many parts of the U.S. Epidemiological studies have suggested that sulfate-related particles are among the most strongly associated with health impacts and premature mortality in adults due to heart attacks, respiratory disease and lung cancer.^{22,23,24}

A recent report estimated that 30,000 premature adult deaths a year occur because of particulate matter.²⁶ However, children may be at even higher risk for particulate matter exposure than adults.²⁷ One factor contributing to this higher risk may be that their exposure to fine particulate matter can be much higher than adults.²⁸ Another factor may be that children are more susceptible to the effects of particulate matter


than adults. Studies in the U.S. have shown that emergency room visits by asthmatic children increase when particulate matter levels rise just slightly above the national air quality standards.^{29,30} Moreover, a Seattle study found that emergency room visits by asthmatic children increased even at fine particulate levels *lower* than EPA's air quality standard.³¹ Some children in the U.S. may be more susceptible to



particulate matter than other children in the population. In one of the first studies of its kind, researchers are evaluating how particulate matter exposure affects African American children with asthma. Results suggest that even small increases in particulate matter may substantially increase asthma symptoms in these children. Results were examined relative to socio-economic factors relating to access of medical care; relationships remained regardless of whether or not their families had contact with a physician for asthma management, other than emergency room visits.³² Seventy-eight percent of African Americans live within 30 miles of a power plant.³³

Figure 2 –
Average annual fine particulate matter conditions in the United States, 1999-2000 (EPA)²⁵

1997 National Ambient Air Quality Standard for fine particles is 15 micrograms per cubic meter of air (ug/m3)



The benefits of reducing particulate matter are clear. For example, in a study undertaken in Germany, changes in respiratory disorders in children were tracked as particulate matter and sulfur dioxide in the air declined in East Germany following the fall of the Berlin Wall.³⁴ The results suggest that non-asthmatic respiratory symptoms such as coughing, chronic bronchitis, ear infections, frequent colds and febrile infections declined in parallel with improving air quality.

Lung Growth in Children is Decreased by Particulate Matter

Can exposure to particulate matter permanently affect a child's developing lungs? The Children's Health Study in California study suggests that particulate matter (PM₁₀) may slow lung function growth in children. Children examined in a dozen communities near Los Angeles experienced a three to five percent relative reduction in lung function *growth* between the most polluted and least polluted cities as a result of exposure to particulate matter.³⁵ When children moved to communities with higher particulate matter, a decreased growth in lung function was observed.³⁶ Conversely, for those children who moved to communities with cleaner air, lung function growth rates increased. This suggests serious permanent harm may befall children living in areas polluted with particulate matter.

Unborn Children are at Risk from Particulate Matter



Particulate matter may affect children even before they are born. Low birth weights and premature births may result due to exposure of a developing fetus to particulate matter in utero. Research in a coal-dominated

region of the Czech Republic indicates that stunted development of the fetus may be a result of exposure to very high levels of particulate matter.³⁷ In fact, in utero exposure to a mixture of power plant pollutants (including sulfur dioxide, nitrogen oxides, polycyclic aromatic hydrocarbons and metals) were associated with neurobehavioral impairment and learning disabilities that extended to 8th grade.³⁸

Similarly, researchers in China have found that high particulate matter concentrations may affect a developing baby. In a study of 75,000 births, an approximate 10 percent higher risk of having a low-birth weight baby associated with increases in sulfur dioxide and total particulate matter levels.³⁹

Particulate Matter Increases the Risk of Infant Death

In the U.S., lung disease and breathing problems are the number one killers of babies under the age of one year, and air pollution is clearly a contributor. Breathing problems accounted for thirty percent of all infant deaths in 1997.⁴⁰ In a comparison of 86 cities in the U.S., researchers found that infants who lived in a highly polluted city during their first two months of life had a mortality rate ten percent higher than infants living in the city with the cleanest air.⁴¹ Investigators in this study found that high particulate matter levels were associated with a 26 percent increased risk of Sudden Infant Death Syndrome and 40 percent increased risk of respiratory mortality.⁴² In a preliminary study extending this work, researchers recently estimated that eleven percent of the infant mortality in the United States is attributable to particulate matter even at low to moderate levels.^{43,44} Studies in the Czech Republic and Mexico City previously had supported the relationship between elevated particulate matter levels and increases in infant mortality.^{45,46}



Ozone Smog Harms Developing Lungs

Ozone, Another Byproduct of Fossil Power is Hazardous to Children

Millions of children live in areas that violate national air quality standards for the ozone (see Table 1). Ground level ozone—the main component of smog—is formed in the presence of sunlight from nitrogen oxides and hydrocarbon vapors emitted by power plants, motor vehicles and industrial processes. Power plants are the source of approximately one quarter of all nitrogen oxide emissions in the U.S. While ozone in the upper levels of the atmosphere provides a protective layer from the sun’s ultraviolet radiation, ozone smog at ground level is extremely harmful to lungs.

The respiratory health effects of ozone have been well documented.⁴⁷ According to EPA, short-term exposure to ozone can cause rapid, shallow breathing and related airway irritation, coughing, wheezing, shortness of breath, and exacerbation of asthma, particularly in sensitive individuals and asthmatic children. Short-term ozone exposure also suppresses the immune system, decreasing the effectiveness of bodily defenses against bacterial infections. In research studies, markers of cell damage increase with ozone exposure. An increase in symptoms means an increase in hospital usage. As shown in Figure 3, ozone is a pervasive problem throughout the eastern half of the United States and California. But ozone is also increasing in the West.

Because air quality is typically the poorest in the summer, studies performed on kids at summer camp provide an excellent setting for examining the effect of air pollution on children. At camp, kids are highly active and are constantly exposed to outdoor air pollution. Moreover, pollution levels can be accurately measured at or near the camps. In typical field studies, children are asked several times a day to perform tests that measure their lung function. The lung function data are then compared to rises and falls in pollution levels. Studies performed in the Northeast, southern California, and Canada show clear decreases in lung function associated with exposure to ozone pollution. In addition, the studies demonstrate that the higher the ozone, the more lung function decreased.⁴⁸



Figure 3 – Counties in the United States exceeding the level of the 8 hour standard in 1999. (U.S.EPA)

Ozone Exposures Result in Pediatric Emergency Room Visits and Hospitalizations

Emergency room visits for asthmatic children are strongly linked to ozone levels. Especially during the summer months, daily hospital admissions and emergency room visits increase as ozone levels increase. These relationships have been demonstrated in the U.S., Mexico and Canada.^{50,51,52, 53} Ozone-related asthma attacks also can result in missed school days. In California, absences from school were correlated with daily changes in ozone.⁵⁴

State	Children	State	Children	State	Children
AL	359,885	KY	413,100	NJ	1,166,298
AR	15,925	LA	451,898	OH	2,076,127
AZ	555,791	MA	927,975	OK	288,184
CA	5,633,990	MD	941,629	PA	2,126,349
CT	723,218	ME	116,439	RI	36,336
DC	117,092	MI	1,281,472	SC	318,427
DE	163,341	MO	351,897	TN	703,083
FL	268,654	MS	81,006	TX	2,528,719
GA	575,431	NC	686,109	VA	554,056
IL	1,491,773	NH	150,169	WI	483,516
IN	702,749	NY	1,483,078	WV	118,273

Table 1 – Number of children by state living in counties that exceed the level of the 8-hour ozone standard.⁴⁹

Can Ozone Result in Premature Death?

For many years, researchers have been investigating the potential association between ozone and premature death. While still hotly debated, there is a growing body of evidence supporting such a relationship for adults.⁵⁵ But can ozone exposure result in premature death in children? One study suggests that exposure to ozone and nitrogen oxides in Mexico City is linked to infant deaths.⁵⁶ Interestingly, the study also found that the relationship between air pollution and infant death was even stronger when particulate matter levels were included in the analysis.

Ozone is Associated with Adverse Birth Outcomes and May Retard Lung Development

Adverse birth outcomes such as premature birth and low birth weight are fairly common in the U.S. Ten percent of all U.S. births are pre-term (more than 3 weeks premature) and some of these premature births may be due to air pollution.⁵⁹ These and other effects on the fetus put children at risk for adverse long-term health problems and recently, researchers have begun to more deeply explore the association of ozone and other pollutants with adverse birth outcomes. For example, a new California study suggests associations between ozone exposures during the second month of gestation and

aortic valve defects, pulmonary artery and valve anomalies and other defects.⁶⁰

Other evidence also suggests that exposure to ozone may permanently affect the lung structure of children. In one of the most stunning new developments, researchers at the University of California suggest that lung development may be stunted by regular exposure to ozone that appears to cause fundamental changes in lung and related brain development. In the study, monkeys exposed to ozone developed little more than half of the normal number of branches of their lungs compared to monkeys exposed only to clean air.⁶¹ The lung receives oxygen from the alveoli at the end of the branches and therefore researchers suspect that fewer branches could cause more difficulty breathing.⁶² Researchers also observed that pollutants caused changes in the brains of these monkeys that made them “hypersensitive” and more likely overreact to irritants. This study provides compelling evidence of the potential serious long-term harm to young children by ozone smog.

Some Children are More Susceptible to Ozone than Others

While scientists have documented that children are generally more susceptible to ozone pollution than adults, asthmatic children are even more vulnerable. However, some subgroups of asthmatic children appear to be more susceptible than others. A recent study suggests that asthmatic children that were born pre-term and/or with low birth weights, are at greater risk from ozone exposures.^{63,64} The same study found that susceptibility varied among inner city Black and Hispanic populations as well. Affluence may play a role in susceptibility; children in homes with air conditioners suffered lower exposures than those in homes without because air conditioners are effective in reducing indoor ozone levels.⁶⁵

Can Ozone Pollution Cause Asthma?

We know that inhaling ozone exacerbates asthma in children, but can it cause asthma? The answer is not clear at this time, but a few new studies point in that direction. Researchers examined associations between exercise and asthma in 3500 children from 12 communities with high ozone levels in Southern California. They found that 265 new cases of asthma were diagnosed in the five years following exposure.⁵⁷ Moreover, children living in high ozone communities and who played 3 or more sports were three times more likely to develop asthma compared to children who didn't play sports. The researchers concluded that new asthma diagnoses were associated with heavy exercise in areas with high ozone levels. This is some of the first evidence that ozone may cause asthma. In another California study, scientists related cumulative lifetime ozone exposures to small airway lung function and found evidence of early indicators of chronic obstructive pulmonary disease.⁵⁸

Sulfur Dioxide Asthmatic Children at Risk

Harmful sulfur dioxide gas directly affects nearby communities

Sulfur dioxide is a dangerous gas that adversely affects human health throughout the U.S. (Figure 4) and can especially affect people living in the shadow of power plant smokestacks where impacts are highest. Power plants are the predominant source of sulfur dioxide, emitting 67 percent the emissions in the U.S. One hundred thousand children with asthma lived in counties in 13 states that violated the sulfur dioxide standard in 1998.^{66,67} Despite improvements resulting from the 1990 Clean Air Act amendments, there were 31 nonattainment areas in the U.S. for sulfur dioxide in 1999.⁶⁸ Unfortunately, serious health effects are also associated with sulfur dioxide levels well below the national ambient air quality standard.⁶⁹ Moreover, acute impacts of sulfur dioxide gas are observed in short-term (less than 5 minute) spikes in concentrations. In fact, in 1996 EPA found the current national ambient air quality standards for SO₂ not to be adequately protective for short-term exposures to asthmatic individuals.⁷⁰ While EPA declined to adopt a new short-term standard, it established guideline for reducing this risk at the state level.

Health Impacts of Sulfur Dioxide Emissions

Sulfur dioxide has been associated with health effects ranging from asthma attacks to premature death. It is an irritant that has been shown in both laboratory and epidemiology studies to exacerbate respiratory disease such as asthma, coughing, wheezing, shortness of breath, and reduce lung function in general.⁷² Inhalation is associated with upper respiratory symptoms including nasal congestion and inflammation. Sulfur dioxide gas can also destabilize normal heart rhythms.⁷³ Controlled laboratory and epidemiology studies have demonstrated that children and people with constrictive pulmonary disease such as asthma are at increased risk from exposure to sulfur dioxide.⁷⁴ Asthmatics in particular can suffer when exposed to sulfur

dioxide. These individuals are commonly stricken with shortness of breath, coughing, wheezing and reductions in lung function. Moreover, sulfur dioxide gas is toxic following only minutes of exposure. Exercising asthmatics can experience lung constriction within 5-10 minutes of exposure.⁷⁵

Sulfur dioxide can be deadly. Numerous studies link sulfur dioxide with bronchial reactions, reduced lung function and premature death.⁷⁶ Some studies associate even very small sulfur dioxide exposures with premature death.⁷⁷ Indeed, sulfur dioxide has also been associated with low birth weight and increased risk of premature death at levels below the national ambient air quality standards. Reproductive effects such as reduced sperm quality have also been linked with sulfur dioxide exposure.⁷⁸ A study of infant mortality in the Czech Republic associated high sulfur dioxide exposures⁷⁹ with a 74 percent higher risk of infant death from respiratory causes.⁸⁰

Sulfur dioxide gas is considered to be most toxic in the presence of ozone and particulate matter, a common mixture that results from power plant emissions.⁸² Prior exposure to ozone has been shown to lead to greater sensitivity to sulfur dioxide in adolescents.

Asthma Cases Have Doubled.
Cases of asthma have rapidly increased, more than doubling in the past two decades. Children age 17 and under account for 29 percent of all the asthma cases in the U.S. and 6 percent of US children have asthma.⁷¹ In 1992, 10 percent of U.S. children with asthma were hospitalized sometime during the year.



Figure 4 –
Areas at Risk for Sulfur Dioxide Exposure (U.S. EPA, 1999).⁸¹

Air Toxics Hazardous to Our Children

Air Toxics and the Environment: The Role of Coal

Coal-fired power plants emit many air toxics. In EPA stack tests, 67 different pollutants have been detected in the flue gas of coal-fired power plants.⁸³ Of these, 55 are known to affect the development of a child's brain or nervous system or to affect the way a child's body develops.⁸⁴ Of these 55, 24 are characterized by EPA as either: known, possible, or probable human carcinogens.⁸⁵ These toxic pollutants are formed during the combustion of coal or are present in coal (e.g., metals like mercury and arsenic), and subsequently released through the smokestack. The amount and type of pollutant emitted from the smokestack depends on the combustion temperature, other characteristics of the coal such as ash content, and the type of pollution control devices in place.

Table 2 summarizes recent emission estimates from the Toxics Release Inventory (TRI) for several important toxics and illustrates the magnitude of power plant emissions relative to other source categories. Because

power plants are currently uncontrolled for air toxics, as other source categories are controlled, power plants will account for a larger and larger share of air toxic emissions nationally. Furthermore, in an analysis of fuel use through the year 2010, the EPA predicts that emissions from coal-fired power plants will increase commensurate with an increased demand for electricity.⁸⁶

Health Effects of Air Toxics

Some air toxics are carcinogens and others are neurotoxins (which affect the development of a child's brain or nervous system) or developmental toxins (which affect the way a child's body develops).⁸⁷ Brain development begins in utero and continues until about age 14.⁸⁸ Normal brain development consists of cell formation and organization that takes place at precise times during gestation in a precise sequence. Interference with any stage of these cellular events may have long-term effects.⁸⁹ The timing, pattern and level of exposure largely determine which parts of the brain will be affected and to what degree.⁹⁰ Thus, exposure of pregnant women and women of childbearing age to air toxics is a

**Table 2 –
1999 Toxics Release Inventory (TRI) Data for Selected
Power Plant Air Toxic Emissions.**

	Total Air Emissions (lbs), TRI 1999	Percent of National TRI Air Emissions	Health Effects
Arsenic and compounds	298,297	48 percent	Neurotoxin / Endocrine disruptor / Known human carcinogen of high potency
Beryllium and compounds	8,585	81 percent	Probable human carcinogen of moderate to low potency
Chromium and compounds	1,053,160	25 percent	Known human carcinogen of high potency (Chromium VI)
Manganese and compounds	3,273,899	16 percent	Neurotoxin
Mercury and compounds	Not reported	33 percent (EPA, 1998)	Methylmercury is a known human neurological and developmental toxin and a possible human carcinogen. Elemental mercury is a neurotoxin. Inorganic mercury can cause kidney damage and is a possible human carcinogen.
Nickel and compounds	1,502,569	50 percent	Neurotoxin nickel refinery dust and nickel subsulfide are known human carcinogens. Nickel carbonyl is a probable human carcinogen.
Selenium and compounds	660,424	77 percent	Neurotoxin selenium sulfide is a probable human carcinogen.
Hydrogen chloride	666,193,000	92 percent	Strong respiratory irritant
Hydrogen fluoride	72,700,182	80 percent	Neurotoxin / Strong respiratory irritant

concern. The potential effects of exposure to these pollutants may be cancer or a range of developmental problems such as learning disabilities, attention deficits, loss of IQ points or other disorders depending on the severity of exposure. Many air toxics are also respiratory irritants that can worsen conditions such as asthma.


Children are Exposed to Power Plant Air Toxics in Many Ways

Some air toxics bioaccumulate in the food chain and are ingested as part of our diet, while still others are dispersed globally and contaminate regions far from the emitting source. Exposure to air toxics from power plants may occur from direct inhalation of air pollutants, or "indirect exposure" including the ingestion of meat, dairy products and fish, as well as water, soil, or vegetation that becomes contaminated by air emissions that have deposited to earth and accumulated in the food chain. Pollutants for which indirect exposure is particularly important are mercury, arsenic, dioxins, cadmium and lead. Some toxics may be absorbed through the skin. Absorption through the skin of some power plant air toxics may also occur, especially from direct contact with contaminated water or soil. Another important exposure pathway for children is the ingestion of contaminated soil during play.

Children also can be exposed to contaminants in power plant combustion wastes. Minefilling (i.e., dumping large volumes of combustion waste in abandoned mines) and the disposal of combustion waste in unlined surface impoundments and landfills can contaminate groundwater (a source of drinking water). Agricultural uses of combustion waste as a soil amendment directly contaminates the soil and can contaminate nearby areas with windblown dust. Combustion waste is largely made up of ash and other uncombusted materials that are left when coal and oil are burned. Each year more than 100 million tons of waste is generated from coal and oil combustion. These wastes contain concentrated levels of numerous contaminants, particularly metals like mercury, arsenic, lead, chromium and cadmium.

How Hazardous Pollutants from Power Plants Get into the Environment

Because power plant waste is generally disposed of at the plant site, children living in the vicinity of power plants experience the highest exposure to coal combustion waste and consequently have the highest risk of adverse health effects. Using computer models, EPA assessed the potential health risks to children if power plant



How hazardous materials from power plants get into the environment.

combustion waste leaked from surface impoundments or ash landfills and contaminated drinking water wells.⁹¹ EPA found that if children drink, over a period of years, an average amount of water contaminated with combustion waste, they will have a higher risk of cancer and other health effects. The EPA found the highest potential risks to children from contaminated groundwater were from arsenic, chromium VI, nickel and selenium.

The EPA also analyzed other ways children might be exposed to power plant combustion waste such as inhalation and the ingestion of fruit, vegetables, beef and dairy products contaminated by the wastes.⁹² The highest risks predicted for inhalation were from chromium VI that is dispersed with dust from uncovered ash landfills. Arsenic, barium, beryllium and posed the highest risks from ingestion of food. In fact, according to EPA analyses, children exposed to power plant combustion wastes could have a considerably higher risk of developing cancer and other illnesses than non-exposed children.⁹³

These children may be even more vulnerable to the health risks posed by these facilities because many of them are living in poverty (see Table 3). Living in poverty is usually associated with poor nutritional status, limited access to health care and substandard housing conditions (including exposure to lead paint), all of which may make these children more susceptible to the effects of toxins in their air and food.

Children at Risk from Air Toxics

Health risks from power plant toxics depend on the severity and duration of the exposure, the exposure pathway, whether the child is especially sensitive to the pollutant, and the pollutant's toxicity. Studies that have attempted to quantify the health risks of toxic emissions are limited because they fail to account for multiple and

cumulative exposure to many pollutants at the same time.⁹⁴ This has resulted in assessments that generally underestimate the health risk from exposure to power plant emissions.

Air of Concern: Mercury Emissions from Power Plants

Of the hazardous air pollutants from power plants, the pollutant of greatest concern is mercury. Two national reports by the EPA have implicated coal-fired power plants as the largest industrial emitters of mercury, producing over one third of all mercury pollution in the U.S.^{96,97}

But, inhaling airborne mercury is not the problem. Airborne mercury eventually deposits in water bodies where it is converted to methylmercury and accumulates in fish tissue. The ingestion of mercury-contaminated fish by

expectant mothers and the subsequent exposure of the developing fetus to methylmercury poses the highest risk to children.

Once emitted, mercury can be transported long distances in the atmosphere. Water bodies are contaminated when mercury in the atmosphere deposits to earth in rain and other forms of precipitation. Run-off and erosion into water bodies is also a source of contamination.⁹⁸ In the aquatic environment, mercury is converted by microorganisms into methylmercury. Fish absorb methylmercury from water as it passes over their gills and as they feed on other aquatic organisms. As larger fish eat smaller ones, concentrations of the pollutant increase in the bigger fish, a process known as



Table 3 – Children living within one mile of a coal-fired power plant.⁹⁵

	U.S. Population (November 2000 Projections)	Within 1 Mile of a Coal-Fired Power Plant
Total U.S. population	276,059,000	836,097
Total children (under age 19)	78,537,000	245,400
Children in poverty	12,845,000	48,477
Childhood poverty rate	14.8 %	19.8 %
Percent non-white	17.8 %	21.5 %

bioaccumulation. Consequently, larger predator fish have higher concentrations as a result of eating contaminated prey.⁹⁹

Mercury contamination in fish across the United States is so pervasive that health departments in 42 states have issued fish consumption advisories.¹⁰⁰ In addition, 11 states have consumption advisories for every inland water body for at least one fish species; 6 states have consumption advisories for canned tuna, and 8 have statewide coastal marine advisories for king mackerel. The U.S. Food and Drug Administration has also issued a consumer advisory for pregnant women, women of child-bearing age, nursing mothers and young children. These groups are advised not to eat swordfish, tilefish, shark and king mackerel because of high mercury levels.¹⁰¹

Methylmercury is both a developmental toxin and a neurotoxin. A spectrum of health effects has been observed following exposure, with the severity of effects depending largely on the amount and timing of exposure.¹⁰² Children and the developing fetus are most vulnerable to mercury exposure. Methylmercury that is consumed by the mother passes through the placenta to the developing fetus. Mercury exposure prior to pregnancy is as critical as exposure during pregnancy because it persists in tissues and is slowly excreted from the body. The first weeks of pregnancy also represent a critical time for fetal development. Women of childbearing age (i.e., 15 to 44 years of age) and pregnant women are therefore the most important members of the population in terms of mercury exposure.¹⁰³

In addition to exposure in utero, infants and children ingest methylmercury from breast milk and other foods in their diet. Children and infants are sensitive to mercury's effects because their nervous systems continue to develop until about age 14. Children also have higher exposures than adults because a child eats more food relative to his or her body weight than an adult does. As a result, they have a higher risk for adverse health effects than adults do.¹⁰⁴

Methylmercury interferes with the development and function of the central nervous system. Prenatal exposure from maternal consumption of fish can cause later impairments in children. Infants appear normal during the first few months of life, but later display subtle effects. These effects include poor performance on neuro-

behavioral tests, particularly on tests of attention, fine motor function, language, visual-spatial abilities (e.g., drawing), and memory. These children will likely have to struggle to keep up in school and might require remedial classes or special education.¹⁰⁵

There is also evidence in humans and animals that exposure to methylmercury can have adverse effects on the developing and adult cardiovascular system, blood pressure regulation and heart-rate variability.¹⁰⁶

The Children Left Behind

On March 2, 2001, the Centers for Disease Control and Prevention (CDC) released preliminary findings on mercury levels in blood and hair from the 1999 National Health and Nutrition Examination Survey (NHANES). This is the first time that human tissues have been systematically analyzed for this mercury. The survey found that 10 percent of the women of childbearing age tested had mercury in their blood at levels above the level that the EPA considers to be safe.^{107,108} Nationally, this translates into 6 million women of childbearing age with elevated levels of mercury from eating contaminated fish, and approximately 390,000 newborns at risk of neurological effects from being exposed in utero to elevated levels of mercury.¹⁰⁹

While power plants are not the only source of mercury, three facts are clear: (1) power plants are the largest emitting source category in the U.S., (2) as other sources are controlled, power plants become a bigger factor; and (3) mercury from power plants have yet to be regulated at the state or national level.

There is also statistical evidence that a number of disorders that have been linked to environmental toxins are increasing. These disorders include premature birth and low birth weight, structural birth defects and behavioral and learning disorders.¹¹⁰ How



much of this increase can be attributed to environmental exposures? The National Academy of Sciences (NAS) recently concluded that as many as 3 percent of known developmental and neurological deficits in children are caused by exposure to known toxic substances, including developmental and neurological toxins. The panel also concluded that 25 percent of these problems may be the result of environmental and genetic factors working in combination, and that toxic substances may play a significant but yet undetermined role.¹¹¹

Using this estimate, the National Environmental Trust (NET), Physicians for Social Responsibility and the Learning Disabilities Association of America calculated that 360,000 children – or 1 in 200 children suffer from developmental or neurological defects caused by exposure to known toxic substances including developmental and neurological toxins.¹¹² They note however that this number is likely underestimated because the NAS considered only known developmental and neurological toxicants. Remarkably, information about potential neurotoxicity and developmental neurotoxicity

is essentially absent even for the chemicals produced in the highest volumes.¹¹³ Also, the NAS estimate of neurological and developmental defects refers only to well-recognized and clinically diagnosed mental and physical disabilities. This underestimates other subtle mental and physical deficits that are difficult to diagnose because of the extended time between exposure and effects, or due to effects that have not yet been recognized.¹¹⁴

Scientists recognize that in addition to environmental toxins, genetics and a child's physical and home environment also contribute to developmental disorders in children. No one can say with certainty to what degree a child's impairment is caused by exposure to any one of these particular factors.¹¹⁵ We do know however, that power plant pollution is a contributing factor and we are certain that exposure to these pollutants can be minimized.



Greenhouse Gas Mitigation Benefits to Children's Health

The health of America's children is linked to climate change too. Scientists in the U.S. and worldwide agree that changes in the Earth's atmosphere are occurring due to the build up of carbon dioxide in our atmosphere from burning fossil fuels. In fact, power plants release 38 percent of all of the carbon dioxide emitted from burning fossil fuels in the U.S.¹¹⁶ By reducing emissions from power plants and other CO₂ sources we can diminish the impact of climate change and with it the expected spread of infectious diseases, increased heat and cold-related illnesses, and increased smog formation and the resultant respiratory illnesses.

In a 2001 report to the President, a panel of the National Academy of Sciences concurs with the Intergovernmental Panel on Climate Change (IPCC) scientific report that concludes that global average surface temperatures may increase by 3-10 degrees Fahrenheit by the end of the 21st Century as a result of human activities.¹¹⁷

The potential health impacts associated with climate change are still being assessed, however, based on what we know some early conclusions can be drawn.^{118,119} Potential human health risks include increased prevalence of infectious disease associated with increasing local temperatures, increased cases of heat related stress and illness, and increased exposure to secondarily-formed pollutants, such as ozone smog, the formation of which is largely dependent on heat and sunlight. Diseases common to tropical and warmer areas could spread; especially those carried by mosquitoes and other insects. These diseases could become progressively more common as warmer temperatures enable these insects to become established further north.

Ozone smog is both a potent greenhouse gas and unhealthy to breathe. As discussed in detail earlier in this report, ozone is associated with increased pediatric asthma, emergency room visits hospitalizations, and lost school days. As the climate gets warmer ozone levels are likely to increase. Atmospheric chemists suggest that warming alone may enhance smog formation; a warming of 4 degrees Fahrenheit could increase ozone concentrations by about 5 percent.¹²⁰

A few studies have attempted to estimate the health benefits of strategies to abate greenhouse gases.^{121,122} The Working Group on Public Health and Fossil Fuel Combustion of the World Health Organization in 1997 projected that "business as usual" energy policies between 2000 and 2020 would lead to 8 million premature deaths from increases in particulate matter alone.¹²³ The same study estimated that 700,000 of these deaths would be avoidable. Another international study recently suggests that many lives could be improved or even saved under greenhouse gas mitigation policies that would result in reduced ozone and particulate matter. According to the study, a 10 percent reduction in ozone and particulate matter as a result of applying climate policies in four cities studied in North and South America would result in avoiding 64,000 premature deaths along with 6.1 million asthma attacks and 37 million work loss days.¹²⁴ While the more heavily polluted South American cities stand to gain the most from greenhouse gas mitigation, the study estimated that 56 neonatal deaths and 3000 pediatric hospitalizations would be avoided each year in New York City alone.

It is apparent that climate strategies that reduce greenhouse gases from power plants and other sources will have a positive effect on human health by decreasing levels of other pollutants such as ozone, lessening the risk of the spread of infectious diseases, and reducing temperature related stress and illness.¹²⁵



What Should be Done? Wise Choices To Improve Children's Health

Air pollution from power plants imposes a serious health burden on our children. We know that pollutants have a greater impact on children than adults. These impacts range from minor coughs to asthma attacks, missed school days, hospitalizations, neurological damage, increased risk of cancer and infant death. The pollutants that affect children are also present in our homes (e.g., particulate matter) and our food (e.g., mercury). While the benefits of reducing power plant pollution have been estimated for adults and are many times the cost of emissions controls, little work has been done to quantify the benefits for children. Certainly the benefits will be great. Saving an infant from death means many decades of life preserved.

Polluting coal-fired power plants must be made to comply with modern emission control standards. In addition, the nation's power fleet should be held to nationwide caps on all four of the key types of power plant pollution including nitrogen oxides, sulfur dioxide, mercury and other air toxics, and carbon dioxide. Reducing power plant emissions of nitrogen oxides and sulfur dioxide by 75 percent or more beyond current legal requirements will dramatically reduce fine

particulate matter pollution so that children can breathe more easily.

The threat of power plant air pollution to children can be reduced comprehensively only when the Clean Air Act's 30-year loophole that allows old, dirty power plants to avoid modern standards is finally closed. Technologies that can reduce sulfur dioxide, nitrogen oxides emissions 90 percent or greater are proven and also appear feasible for mercury. The threat posed by global warming requires that we address carbon dioxide as part of a comprehensive strategy to address power plant pollution. Requirements such as these can ensure that U.S. energy policy better accounts for public health and the environmental costs associated with electricity production and will propel us toward a more sustainable energy future that relies increasingly on cleaner sources of energy including renewable energy resources and conservation. Our children's health and quality of life are harmed by air pollution today. The specter of global warming hangs over their future. We can leave our children a legacy of

cleaner air and an improved environment by making wise choices today. Let's make comprehensive power plant clean up our gift to them.



Appendix A: State Statistics on Power Plants and Children.¹²⁶

Many schools are located within the reach of toxic power plant emissions.

State	Major power plants in state	# Power plants within 30 miles of state border	# Children in 30 mile radius of a power plant	# Children in state with asthma	# Schools in 30 mile radius of a power plant
AK	2	2	641	11,567	0
AL	9	15	571,681	66,899	1,992
AR	3	5	203,819	40,955	841
AZ	6	7	186,909	75,689	232
CA	0	1	1,737	551,293	1
CO	14	18	735,646	61,730	713
CT	1	5	441,884	48,971	600
DC	0	5	117,092	6,781	389
DE	3	7	138,477	10,976	266
FL	12	13	1,316,810	211,552	1,723
GA	12	18	1,148,023	121,251	2,379
HI	–	–	–	18,865	–
IA	19	24	484,162	44,879	1,016
ID	0	1	5	21,567	0
IL	27	57	2,688,974	195,748	7,464
IN	29	53	1,102,394	92,614	1,743
KS	8	15	292,989	42,571	1,051
KY	20	44	731,359	60,140	1,326
LA	3	6	204,214	75,539	568
MA	4	6	1,041,591	89,264	1,729
MD	6	19	1,078,810	78,564	2,075
ME	0	2	25,116	18,549	17
MI	20	24	1,769,955	157,438	3,079
MN	14	20	836,183	77,082	1,147
MO	20	35	1,079,199	86,509	4,630
MS	3	7	167,482	46,884	537
MT	2	3	37,693	14,315	163
NC	14	19	1,040,306	113,747	1,659
ND	8	10	46,353	10,358	172
NE	7	11	292,690	27,407	822
NH	2	3	206,909	18,235	190
NJ	4	12	1,679,559	122,805	1,421
NM	4	8	37,498	30,866	98
NV	3	3	5,343	26,018	12
NY	13	16	3,365,226	281,701	3,453
OH	27	46	2,487,909	176,271	7,107
OK	5	9	248,556	54,316	348
OR	1	1	5,811	49,852	12
PA	23	44	2,493,882	178,255	5,117
RI	0	2	212,346	14,650	64
SC	13	20	815,774	58,863	3,885
SD	2	5	35,333	12,552	159
TN	7	14	757,459	81,754	2,817
TX	18	20	1,316,753	339,660	1,059
UT	5	8	22,753	42,338	48
VA	10	20	1,176,637	100,484	2,136
VT	–	–	–	9,027	–
WA	2	2	512,411	88,961	894
WI	16	25	983,789	83,330	1,832
WV	13	27	345,602	26,053	2,987
WY	8	9	58,684	8,260	190
All States	442	746	34,550,428	4,283,955	72,163

MSB Energy Associates / Source: Clean Air Network database.

Appendix B: Ranked Exposure Profiles

Profiles for populations within 30 miles of coal-fired power plants.

State	Children w/ Asthma	Rank	Total Air Releases (tons)	Tons per Child	Air Releases (tons)	No. of Schools	Ranks	No. Children in Non-Attainment Areas
AK	406	47	0	0	44	0	48	0
AL	34,733	21	88,323,224	125	8	1,992	13	189,779
AR	14,090	30	28,644,100	110	29	841	26	0
AZ	13,952	31	47,135,216	201	17	232	35	188,021
CA	339	48	0	0	44	1	47	0
CO	50,336	17	40,535,866	50	19	713	28	524,240
CT	8,715	34	2,287,185	15	43	600	29	151,632
DC	5,244	36	0	0	44	389	32	0
DE	8,312	35	5,670,211	45	39	266	34	115,011
FL	90,859	8	77,702,956	58	10	1,723	17	0
GA	78,897	9	83,778,288	62	9	2,379	10	674,716
IA	30,470	22	40,512,642	70	20	1,016	24	0
ID	71	49	0	0	44	0	48	0
IL	157,659	2	92,931,607	33	6	7,464	1	1,906,193
IN	62,469	14	137,162,285	123	3	1,743	15	208,281
KS	18,157	25	39,717,525	109	22	1,051	23	0
KY	44,158	20	104,420,412	123	5	1,326	20	198,011
LA	17,199	27	24,287,341	68	31	568	30	121,999
MA	28,266	23	13,806,340	30	36	1,729	16	157,453
MD	66,360	12	32,503,474	0	28	2,075	12	800,609
ME	1,541	44	0	0	44	17	44	0
MI	106,194	5	75,587,293	41	11	3,079	7	0
MN	52,479	16	38,300,886	40	24	1,147	21	149,632
MO	65,728	13	71,722,251	63	13	4,630	4	73,998
MS	14,468	29	15,678,705	55	35	537	31	0
MT	2,553	41	17,955,967	417	34	163	39	34,708
NC	74,947	10	73,944,757	63	12	1,659	18	0
ND	1,943	43	37,382,883	996	25	172	38	0
NE	17,706	26	21,435,881	70	32	822	27	111,898
NH	12,630	32	4,706,364	19	40	190	37	86,198
NJ	99,488	6	9,815,066	5	38	1,421	19	1,577,955
NM	3,539	39	33,799,488	541	27	98	41	0
NV	929	45	20,289,092	1,645	33	12	45	5,454
NY	179,051	1	25,760,999	9	30	3,453	6	2,202,581
OH	139,029	4	137,570,633	54	2	7,107	2	479,445
OK	16,611	28	38,849,383	126	23	348	33	0
OR	761	46	4,021,645	316	42	12	46	0
PA	140,949	3	112,280,946	46	4	5,117	3	1,560,240
RI	11,966	33	0	0	44	64	42	0
SC	48,567	19	41,015,630	41	18	3,885	5	0
SD	2,327	42	4,159,081	82	41	159	40	0
TN	49,941	18	65,294,367	68	14	2,817	9	226,307
TX	92,386	7	163,298,619	107	1	1,059	22	682,338
UT	2,603	40	37,118,861	822	26	48	43	3,576
VA	73,891	11	39,780,790	30	21	2,136	11	350,362
VT	0	50	0	0	44	0	48	0
WA	5,116	37	10,451,134	142	37	894	25	56,376
WI	60,829	15	51,081,291	46	16	1,832	14	512,883
WV	19,642	24	92,265,497	235	7	2,987	8	14,165
WY	3,880	38	51,357,490	723	15	190	36	0
U.S. Total	1,968,865		2,154,343,671	57		69,097		12,986,262

Data: MSB Energy Associates

Endnotes

- 1 Cifuentes, L., Borja-Aburto, V., Gouveia, N., Thurston, G. D., and Davis, D. 2001. Assessing the health benefits of urban air pollution reductions associated with climate change mitigation (2000-2020). Santiago, Sao Paulo, Mexico City and New York City. *Environmental Health Perspectives*, vol. 109, supplement 3, p. 419-425, June .
- 2 Clean Air Task Force. 2001. *Cradle to Grave: The environmental impacts from coal*. Clean Air Task Force 77 Summer Street, Boston, MA 02110.
- 3 Bascom, R. et al. 1996. Health effects of outdoor air pollution, Part 1. *American Journal of Respiratory and Critical Care Medicine* vol. 153, pp. 3-50.
- 4 Williams, L., Spence, A. and Tideman, S. 1977. Implications of the Observed Effect of Air Pollution on Birth Weight. *Social Biology*, vol. 24, no. 1, p. 1-9.
- 5 Bascom, R. et al. 1996. Health effects of outdoor air pollution, Part 2. *American Journal of Respiratory and Critical Care Medicine* vol. 153, pp. 477-498.
- 6 Frischer, T., Studnicka, M., Gartner, C., Tauber, E., Horak, F., Veiter, A., Spengler, J., Kühr, J., and Urbanek, R. 1999. Lung Function Growth and Ambient Ozone: A Three-Year Population Study in School Children. *Am J Respir Crit Care Med*, vol. 160, pp. 390-396.
- 7 McCormick, M.C. 1985. The contribution of low birth weight to infant mortality and childhood morbidity. *New England Journal of Medicine* vol. 312 p. 164-167.
- 8 Woodruff, T., Grillo, J. and Schoendorf, K. 1997. The relationship between selected causes of postneonatal infant mortality and particulate air pollution in the United States. *Environmental Health Perspectives*, vol. 105, 608-612.
- 9 Bobak, M., and Leon, D. 1999. The effect of air pollution on infant mortality appears specific for respiratory causes in the postneonatal period. *Epidemiology*. vol. 10, no. 6 pp. 666-670.
- 10 Srivastava, Ravi, K. (2000) *Controlling SO2 Emissions: A Review of Technologies*. Prepared for U.S. EPA, Office of Research and Development; EPA/600/R-00/093 November 2000
- 11 Wiley, J.A., Robinson, J.P., Cheng, Y.T, Piazza, T, Stork, L, Pladsen, K. 1991. Study of children's activity patterns. Final report contract No. A733-149; Survey Research Center, University of California, Berkeley, September 1991.
- 12 Snodgrass, W.R. Physiological and biochemical differences between children and adults and determinants of toxic response to environmental pollutants. In: Guzman, et al., *Similarities and Differences Between Children and Adults: Implications for Risk Assessment*. 1151 Press, Washington, DC. (year unknown).
- 13 EPA Green Book; compiled by MSB Energy Associates;
- 14 Raizenne, M., Dales, R. and Burnett, R. 1998. Air pollution exposures and children's health. *Canadian Journal of Public Health* vol. 89 supplement 1, May-June 1998.
- 15 30 miles was chosen as the metric as it is a standard criteria used in modeling impacts from power plumes using ISC; supporting this is a recent study (Levy et. al., 2000) suggesting that impacts from power plants are greatest within a 30 mile radius of a smokestack in MA.
- 16 U.S. EPA (2001); *National Air Quality and Emissions Trends Report, 1999*. EPA publication 454/R01-004, March 2001. <http://www.epa.gov/airtrends/>
- 17 U.S. EPA.1998. Study of hazardous air pollutant emissions from electric utility steam generating units – final report to Congress. February. 453/R-98-004a.
- 18 From EPA Emissions of Greenhouse Gases in the U.S. 2000: Electric utilities release 2453 million tons per year; total combustion related sources of CO2 total 6386 million tons. Ratio of electric utility CO2 to total combustion related sources is 38%. Data provided by MSB Energy Associates.
- 19 The Potomac River Plant, VA is a 1940s and 1950s vintage power plant next to an apartment complex. The height of the stack is approximately the height of building.
- 20 U.S. EPA, 2000. *National air quality and emissions trends report*. 1998. EPA 454/R-00-003, March.
- 21 US EPA Toxics Release Inventory (TRI) <http://www.epa.gov/tri/tri99/index.htm>
- 22 Pope, C.A., Burnett, R.T., Thun, M.J., Calle, E.E., Krewski, D., Ito, K., Thurston, G.D. (2002). Lung cancer, cardiopulmonary mortality and long-term exposure to fine particulate air pollution. *JAMA*, v. 287, No. 9, p. 1132-1141.
- 23 Thurston, G.D. 1998. Determining the pollution sources associated with PM health effects. *AWMA VIP-81*, v.2 , p 889.
- 24 Dockery, D.W. et. al. 1993. An association between air pollution and mortality in six U.S. cities. *New England Journal of Medicine*, vol. 329 p. 1753-1759.
- 25 Map: MSB Energy Associates based on unpublished EPA data.
- 26 Abt Associates. 2000. *The particulate-related health benefits of reducing power emissions*; Abt Associates, Bethesda, MD.
- 27 Thurston, G. D., 2000. Particulate matter and sulfate: Evaluation of current California air quality standards with respect to protection of children: California Air Resources Board, Office of Environmental Health Hazard Assessment; September 1, 2000. <http://www.arb.ca.gov/ch/ceh/airstandards.htm>
- 28 Brown, K.H., Suh, H.H. and Koutrakis, P. 2001. Characterization of personal-ambient PM 2.5 relationships for children and older adults. *Health Effects Institute Annual Conference, Program and Abstracts*. April.
- 29 Pope, C.A., Dockery, D.W. 1992. Acute health effects of PM 10 pollution symptomatic and asymptomatic children. *American Review of Respiratory Disease* vol. 145, p. 1123-1128.
- 30 Tolbert, P., et al. 2000. Air quality and pediatric emergency room visits for asthma in Atlanta, Georgia. *American Journal of Epidemiology*, vol. 151, no. 8 p. 798-810 .
- 31 Norris, G., Young Pong, N., Koenig, J., Larson, T., Sheppard, L. and Stout, J. 1999. An association between fine particles and asthma emergency department visits for children in Seattle. *Environmental Health Perspectives*, vol. 107, no. 6, pp. 489-493.
- 32 Ostro, B., Lipsett, M., Mann, J., Braxton-Owens, H., White, M. 2001. Air pollution and exacerbation of asthma in African-American Children in Los Angeles. *Epidemiology* vol. 12, no. 2, p. 200-208.
- 33 Based on 2000 data, the number of African Americans living within 30 miles of a power plant was 26.5 million out of a total of 33.9 million in the U.S. population. Thus, 78% percent of all African Americans live within 30 miles of a power plant. Estimated using total U.S. population from the 2000 census and racial fractions from 1990 census data. Data provided by MSB Energy Associates.
- 34 Heinrich, J, Hoelscher, B and Wichmann, HE (2000) Decline of ambient air pollution and respiratory symptoms in children. *American Journal of Respiratory and Critical Care Medicine*, vol. 161, pp. 1930-1936.

- 35 Gauderman, W.J., McConnell, R., Gilliland, F., London, S., Thomas, D., Avol, E., Vora, H., Berhane, K., Rappaport, E., Lurmann, F., Margolis, H.G., and Peters, J. 2000. Association between air pollution and lung function growth in Southern California children. *American Journal of Respiratory and Critical Care Medicine*, vol. 162, no. 4, pp. 1-8.
- 36 Avol, E.L., Guaderman, W.J., Tan S.M., London, S.J., and Peters, J.M. (2001). Respiratory effects of relocating to areas of differing air pollution levels. *American Journal of Respiratory and Critical Care Medicine* v. 164 p. 2067-2072.
- 37 Lewtas, J., Binkova, B., Misova, I., Subrt, P., Lenicek, J. and Sram, R. J. 2001. Biomarkers of exposure to particulate air pollution in the Czech Republic; in Teplice Program: Impact of Air Pollution on Human Health; Academia Press, Prague; ISBN 80-200-0876-4.
- 38 Otto, D., Skalik, I., Hudnesll, K., House, D. and Sram, R. J.2001. Neurobehavioral effects of exposure to environmental pollutants in Czech children; in Teplice Program: Impact of Air Pollution on Human Health; Academia Press, Prague; ISBN 80-200-0876-4.
- 39 Wang, X., Ding, H., Ryan, L., and Xu, X. 1997. Association between air pollution and low birth weight; a community-based study. *Environmental Health Perspectives* vol. 15 p.514-520.
- 40 American Lung Association. 2000. <http://www.lungusa.org/pub/1dd.html>
- 41 Woodruff, T., Grillo, J. and Schoendorf, K. 1997. The relationship between selected causes of postneonatal infant mortality and particulate air pollution in the United States. *Environmental Health Perspectives*, vol. 105, p. 608-612.
- 42 Industry called this study into question citing exposure misclassification due to insufficient assessment of geography/ location and related PM exposures. However, the HEI reanalysis of the American Cancer Society suggests that geographic exposure differences make relatively little difference in PM dose response.
- 43 Kaiser, R. Kunzli, N., Schwartz, J. 2001. The impact of PM 10 on infant mortality in 8 U.S. cities. Abstract, American Thoracic Society, Abstract Preview: ATS1P1_6266.
- 44 PM 2.5 levels over 7.5 ug/m 3.
- 45 Bobak, M. and Leon, D. 1999. The effect of air pollution on infant mortality appears specific for respiratory causes in the postneonatal period. *Epidemiology*. vol. 10, no. 6 p. 666-670.
- 46 Loomis, D., Castillejos, M., Gold, D., McDonnell, W. Borja-Aburto, V. 1999. Air pollution and infant mortality in Mexico City. *Epidemiology*, vol. 10, p. 118-123.
- 47 U.S. EPA. 1996. Air quality criteria for ozone and related photochemical oxidants; EPA/600/P-93/004aF.
- 48 Galizia, A., Kinney, P.L. (1999). Long-term residence in areas of high ozone: associations with respiratory health in a nationwide sampling of nonsmoking young adults. *Environmental Health Perspectives*, v. 107, no. 8.
- 49 MSB Energy Associates
- 50 Thurston, G.D. and Ito, K. 1999. Epidemiological studies of ozone exposure effects. In *Air Pollution and Health*, Stephen T. Holgate et. al., Ed., Academic Press, London.
- 51 Tolbert, P. E., et. al. 2000. Air quality and pediatric emergency room visits for asthma in Atlanta, Georgia. *American Journal of Epidemiology*, vol. 151, no. 8 p. 798-810 .
- 52 White, M.C., Etzel, R.A., Wilcox, W.D. and Lloyd, C. 1994. Exacerbation of childhood asthma and ozone pollution in Atlanta. *Environmental Research* vol. 65, p. 56-68.
- 53 Burnett, R., et. al. 2001. Association between ozone and hospitalization for acute respiratory diseases in children less than 2 years of age. *American Journal of Epidemiology*, vol. 153, no. 5, p. 444-452.
- 54 Peters, J.M., Avol, E., Berhane, K., Gauderman, W.J., Gilliland, F., London, S., Lurmann, H., Margolis, H., McConnell, R., and Thomas, D.C. 2001. Chronic respiratory effects of air pollution in southern California Children; Poster; HEI Annual Conference, Washington DC; Program and Abstracts; Health Effects Institute, Cambridge, MA 02139.
- 55 Thurston, G.D. and Ito, K. 1999. Epidemiological studies of ozone exposure effects. In *Air Pollution and Health*, Stephen T. Holgate et. al., Ed., Academic Press, London
- 56 Loomis, D., Castillejos, M., Gold, D., McDonnell, W. and Borja-Aburto, V.1999. Air pollution and infant mortality in Mexico City. *Epidemiology*. vol. 10, no. 2, p. 118-123.
- 57 McConnell, R., Berhane, F., Gilliland, T., Islam, S.J., Gauderman, W.J., Avol, E., Margolis, H.G., Peters, J.M. 2001. American Thoracic Society abstract (poster) May 20 2001 <http://www.abstracts-on-line.com/abstracts/ATS>.
- 58 Kunzli, N., Lurmann, F., Segal, M., Ngo, L., Balmes, J., and Tager, I.B., 1997. Association between Lifetime Ambient Ozone Exposure and Pulmonary Function in College Freshmen Results of a Pilot Study. *Environmental Research*, Vol. 72, pp. 8-23.
- 59 Ritz, B., (2001) Overview: Epidemiologic studies of air pollution and adverse birth outcomes. Presentation slides from HEI Annual Meeting , May 2001
- 60 Ritz, B., Yu, F., Fruin, S, Chapa, G., Shaw, G., Harris, J (2002) Ambient air pollution and risk of birth defects in southern California . *American Journal of Epidemiology*, v. 155, no. 1.
- 61 Plopper, C.G., Fanucci, M.V., Evans, M.J., Larson, S.P., Schelegle, E.S., Joad, J.P., Pinkerton, K.E., VanWinkle, L.S., Gershwin, L.J., Miller, L.A., Wu, R., Buckpitt, A.R., and Hyde, D.M. 2001. Air pollution effects in a primate model of asthma. Abstract and presentation, HEI Annual Conference, Washington DC; Program and Abstracts; Health Effects Institute, Cambridge MA, 02139
- 62 Edie Lau, Sacramento Bee, April 15, 2001 "Study suggests asthma culprit: young lungs exposed to ozone seem more prone to problems with development.
- 63 Mortimer, K.M., Tager, I.B., Dockery, D.W. Neas, L.M., Redline S. 2000. The effect of ozone on inner city children with asthma. Identification of susceptible subgroups. *American Journal of Respiratory and Critical Care Medicine*, vol. 162, p. 1838-1845.
- 64 The average ozone daily level during the study was 48 ppb.
- 65 Mortimer, K.M., Tager, I.B., Dockery, D.W. Neas, L.M., Redline S. 2000. The effect of ozone on inner city children with asthma. Identification of susceptible subgroups. *American Journal of Respiratory and Critical Care Medicine*, vol. 162, page 1843.
- 66 MSB energy Associates; data from EPA Green Book.
- 67 MSB energy Associates; data from EPA Green Book.
- 68 U.S. EPA (2001); National Air Quality and Emissions Trends Report, 1999. EPA publication 454/R01-004, March 2001. <http://www.epa.gov/airtrends/>

- 69 The National Ambient Air Quality Standards for sulfur dioxide are 140 ppb (24 hour average) or 30 ppb (annual average). Although advocates have argued for a short term NAAQS such as California's short-term standard (1 hour is 250 ppb) or a 5 minute standard, the federal government has not implemented a more protective short-term sulfur dioxide NAAQS.
- 70 U.S. EPA 40CFR Part 50 AD-FRL-National Ambient Air Quality Standards for Sulfur Oxides (Sulfur Dioxide)–Final Decision. May 15, 1996. P. 50.
- 71 MSB Energy Associates. Source: CDC/American Lung Association.
- 72 Koenig, J. Q., Mar, T. F. 2000. Sulfur dioxide: Evaluation of current California air quality standards with respect to children's health; California Air Resources Board, California Office of Environmental Health Hazard Assessment. <http://www.arb.ca.gov/ch/ceh/airstandards.htm>
- 73 Peters, A. et. al. 1999. Increases in heart rate variability during an airpollution episode. *American Journal of Epidemiology*, 150 (10).
- 74 Koenig, J. Q., Mar, T. F. 2000. Sulfur dioxide: Evaluation of Current California Air Quality Standards with Respect to Protection of Children; CARB California Office of Environmental Health Assessment, September.
- 75 U.S. EPA. 1994. Supplement to the Second Addendum to the Air Quality Criteria of Particulate Matter and Sulfur Oxides (1982): Assessment of new findings on sulfur dioxide acute exposure health effects in asthmatic individuals. EPA-600R-93/002; August.
- 76 Koenig, J. Q., Mar, T. F. 2000. Sulfur dioxide: Evaluation of Current California Air Quality Standards with Respect to Protection of Children; CARB California Office of Environmental Health Assessment, September.
- 77 Kelsall, J.E., Sarney, J.M., Zeger, S.L., Ku, J. 1997. Air pollution and mortality in Philadelphia 1974-1988. *American Journal of Epidemiology*, vol.146, p. 750-762.
- 78 Perrault, S.D., Selevan, S.G., Evenson, D.P. and Rubes, J. 2001. Health Effects Institute Annual Conference April 2001 Program and Abstracts p.28.
- 79 Sulfur dioxide exposures = 50 ug/m3.
- 80 Bobak, M., Leon, D. 1999. The effect of air pollution on infant mortality appears specific for respiratory causes in the postneonatal period. *Epidemiology*. vol. 10, no. 6 p. 666-670.
- 81 Second highest annual maximum 24 hour sulfur dioxide concentration (1999 EPA Air Quality and Emission Trends Report).
- 82 American Thoracic Society.1996. State of the Art. Health Effects of Outdoor Air Pollution. *American Journal of Respiratory and Critical Care Medicine*. vol. 153, no. 1 p.3-50.
- 83 U.S. EPA. 1998. Study of hazardous air pollutant emissions from electric utility steam generating units – final report to Congress. February. 453/R-98-004a.
- 84 National Environmental Trust (NET), et. al. 2000. Polluting Our Future: Chemical Pollution in the U.S. that Affects Child Development and Learning. September. www.environment.org
- 85 U.S. EPA. 1998. Study of hazardous air pollutant emissions from electric utility steam generating units – final report to Congress. February. 453/R-98-004a.
- 86 U.S. EPA. 1998. Study of hazardous air pollutant emissions from electric utility steam generating units – final report to Congress. February. 453/R-98-004a.
- 87 National Environmental Trust (NET), et. al. 2000. Polluting Our Future: Chemical Pollution in the U.S. that Affects Child Development and Learning. September. www.environment.org.
- 88 U.S. EPA. 1997. Mercury Study Report to Congress: Volume I Executive Summary. December. EPA 452/R-97-003.
- 89 Greater Boston Physicians for Social Responsibility (GBPSR). 2001. In Harm's Way: Toxic Threats to Child Development. January . www.igc.org/psr/
- 90 Greater Boston Physicians for Social Responsibility (GBPSR). 2001. In Harm's Way: Toxic Threats to Child Development. January . www.igc.org/psr/
- 91 U.S. EPA. 1998b. Technical background document for the supplemental report to Congress on remaining fossil fuel combustion wastes. Groundwater pathway human health risk assessment. Revised draft final. Office of Solid Waste. June.
- 92 Research Triangle Institute. 1998. Draft final report. Non-groundwater pathways, human health and ecological risk analysis for fossil fuel combustion phase 2 (FFC2). Prepared for the U.S. EPA Office of Solid Waste, Washington, D.C. June 5, 1998.
- 93 U.S. EPA. 1998. Study of hazardous air pollutant emissions from electric utility steam generating units – final report to Congress. February. 453/R-98-004a.
- 94 U.S. EPA. 1998. Study of hazardous air pollutant emissions from electric utility steam generating units – final report to Congress. February. 453/R-98-004a.
- 95 Resident Population Estimates of the United States by Sex, Race and Hispanic Origin: April 1, 1990 to July 1, 1999 with Short-term Projections to November 1, 2000. Public release date: January 2, 2001.
- 96 U.S. EPA.1998. Study of hazardous air pollutant emissions from electric utility steam generating units – final report to Congress. February. 453/R-98-004a.
- 97 U.S. EPA.1997b. Mercury Study Report to Congress. Volume II: An Inventory of Anthropogenic Mercury Sources in the United States. December.
- 98 U.S. EPA. 1997d. Mercury Study Report to Congress, Volume III: Fate and Transport of Mercury in the Environment. EPA-452/R-97-005.
- 99 U.S. EPA. 1998. Study of hazardous air pollutant emissions from electric utility steam generating units – final report to Congress. February. 453/R-98-004a.
- 100 <http://www.epa.gov/ost/fish>
- 101 FDA Consumer Advisory for Pregnant Women and Women of Childbearing Age who may become Pregnant about the Risks of Mercury in Fish. March 2001. <http://www.cfsan.fda.gov/~dms/qa-pes1.html>
- 102 U.S. EPA. 1997f. Mercury Study Report to Congress, Volume V: Health Effects of Mercury and Mercury Compounds. EPA-452/R-97-007.
- 103 U.S. EPA. 1997b. Mercury Study Report to Congress, Volume VII: Characterization of Human and Wildlife Risks from Mercury Exposure in the United States. EPA-452/R-97-009
- 104 U.S. EPA. 1997b. Mercury Study Report to Congress, Volume VIII: Characterization of Human and Wildlife Risks from Mercury Exposure in the United States. EPA-452/R-97-009

- 105 Toxicological Effects of Methylmercury. National Academy Press, Washington, DC, 2000. <http://www.nap.edu>
- 106 Toxicological Effects of Methylmercury. National Academy Press, Washington, DC, 2000.
- 107 U.S. Centers for Disease Control and Prevention. 2001. Blood and hair mercury levels in young children and women of childbearing age - United States, 1999. Morbidity and Mortality Weekly, March 2, 2001.
- 108 The EPA has estimated how much methylmercury is harmful to the developing fetus by developing a "reference dose" or RfD. An RfD is a dose that can be ingested daily over a lifetime without harmful health effects.
- 109 Derived from 1990 census data. <http://www.census.gov>
- 110 National Environmental Trust (NET), et. al. 2000. Polluting Our Future: Chemical Pollution in the U.S. that Affects Child Development and Learning. September. www.envirnet.org
- 111 National Academy of Sciences (NAS). 2000. Scientific Frontiers in Developmental Toxicology and Risk Assessment. National Academy Press, June. <http://www.nap.edu/books/0309070864/html/>
- 112 National Environmental Trust (NET), et. al. 2000. Polluting Our Future: Chemical Pollution in the U.S. that Affects Child Development and Learning. September. www.envirnet.org. The Census Bureau estimates that nearly 12 million U.S. children under 18 (17 percent of children) suffer from one or more developmental, learning, or behavioral disabilities. If, according to the National Academy of Sciences, known toxic exposures are directly implicated in approximately 3 percent of these disabilities, then 360,000 U.S. children - or 1 in every 200 children - children suffer from developmental or neurological defects caused by exposure to known toxic substances including developmental and neurological toxins.
- 113 Greater Boston Physicians for Social Responsibility (GBPSR). 2001. In Harm's Way: Toxic Threats to Child Development. January. www.igc.org/psr/
- 114 National Environmental Trust (NET), et. al. 2000. Polluting Our Future: Chemical Pollution in the U.S. that Affects Child Development and Learning. September. www.envirnet.org
- 115 Greater Boston Physicians for Social Responsibility (GBPSR). 2001. In Harm's Way: Toxic Threats to Child Development. January. www.igc.org/psr/
- 116 From EPA Emissions of Greenhouse Gases in the U.S. 2000: Electric utilities release 2453 million tons per year; total combustion related sources of CO2 total 6386 million tons. Ratio of electric utility CO2 to total combustion related sources is 38%. Data provided by MSB Energy Associates.
- 117 National Research Council. 2001. Climate change science. National Academy Press, Washington D.C. ISBN 0-309-07574-2.
- 118 IPCC. 2001. Climate Change 2001: Impacts, adaptation and vulnerability; Summary for Policymakers. <http://www.ipcc.ch/pub/wg2SPMfinal.pdf>
- 119 U.S. EPA. 2001. Global warming impacts summary. <http://www.epa.gov/globalwarming/impacts/health/index.html>
- 120 Hansen, J. 2000. Global warming in the 21st century: an alternative scenario. Proceedings of the National Academy of Sciences, Aug 29, Early Edition.
- 121 Abt Associates.(1999. Co-control benefits of greenhouse gas control policies. Prepared for U.S. EPA. Bethesda, MD.
- 122 Cifuentes, L., Borja-Aburto, V., Gouveia, N., Thurston, G. D., and Davis, D. 2001. Assessing the health benefits of urban air pollution reductions associated with climate change mitigation (2000-2020). Santiago, Sao Paulo, Mexico City and New York City. Environmental Health Perspectives, vol. 109, supplement 3, p. 419-425, June.
- 123 Working Group on Public Health and Fossil Fuel Combustion. Short-term improvements in public health from global climate policies on fossil fuel combustion. Lancet (1997) 350 1341-1349.
- 124 The four cities are New York City, Sao Paulo, Brazil, Mexico City, Mexico and Santiago de Chile.
- 125 National Research Council. 2001. Climate change science. National Academy Press, Washington D.C. ISBN 0-309-07574-2, p. 20.
- 126 MSB Energy Associates. Source: Clean Air Network database.



TERRILYN BAYNE



Clear the Air: National Campaign Against Dirty Power

1200 18th Street N.W.
Washington D.C. 20036
Tel: 202-887-1715
<http://www.cleartheair.org>

Physicians for Social Responsibility

1875 Connecticut Avenue, NW, Suite 1012
Washington DC 20009
Tel: 202-667-4260
<http://www.psr.org>

Clean Air Task Force

77 Summer Street,
Boston, MA 02110
Tel: 617-292-0234
<http://clnatf.org>